



Underuse of Prevention and Lifestyle Counseling in Patients With Peripheral Artery Disease

Jeffrey S. Berger, MD, MS,^a Joseph A. Ladapo, MD, PhD^b

ABSTRACT

BACKGROUND Little is known about patterns of medication use and lifestyle counseling in patients with peripheral artery disease (PAD) in the United States.

OBJECTIVES The authors sought to evaluate trends in both medical therapy and lifestyle counseling for PAD patients in the United States from 2005 through 2012.

METHODS Data from 1,982 outpatient visits among patients with PAD were obtained from the National Ambulatory Medical Care Survey and National Hospital Ambulatory Medical Care Survey, a nationally representative assessment of office-based and hospital outpatient department practice. Trends in the proportion of visits with medication use (antiplatelet therapy, statins, angiotensin-converting enzyme inhibitors [ACEIs] or angiotensin receptor blockers [ARBs], and cilostazol) and lifestyle counseling (exercise or diet counseling and smoking cessation) were evaluated.

RESULTS Over the 8-year period, the average annual number of ambulatory visits in the United States for PAD was 3,883,665. Across all visits, mean age was 69.2 years, 51.8% were female, and 56.6% were non-Hispanic white. Comorbid coronary artery disease (CAD) was present in 24.3% of visits. Medication use for cardiovascular prevention and symptoms of claudication was low: any antiplatelet therapy in 35.7% (standard error [SE]: 2.7%), statin in 33.1% (SE: 2.4%), ACEI/ARB in 28.4% (SE: 2.0%), and cilostazol in 4.7% (SE: 1.0%) of visits. Exercise or diet counseling was used in 22% (SE: 2.3%) of visits. Among current smokers with PAD, smoking cessation counseling or medication was used in 35.8% (SE: 4.6%) of visits. There was no significant change in medication use or lifestyle counseling over time. Compared with visits for patients with PAD alone, comorbid PAD and CAD were more likely to be prescribed antiplatelet therapy (odds ratio [OR]: 2.6; 95% confidence interval [CI]: 1.8 to 3.9), statins (OR: 2.6; 95% CI: 1.8 to 3.9), ACEI/ARB (OR: 2.6; 95% CI: 1.8 to 3.9), and smoking cessation counseling (OR: 4.4; 95% CI: 2.0 to 9.6).

CONCLUSIONS The use of guideline-recommended therapies in patients with PAD was much lower than expected, which highlights an opportunity to improve the quality of care in these high-risk patients. (J Am Coll Cardiol 2017;69:2293-300)
© 2017 by the American College of Cardiology Foundation.

Peripheral artery disease (PAD) is a highly prevalent atherosclerotic syndrome with an estimated global population burden of ~200 million people (1). Because of the aging population and risk factor trends, the prevalence of PAD is increasing (2,3). Patients with PAD are at heightened risk for adverse cardiovascular and limb events, and impaired quality of life (4). In fact, PAD is considered



Listen to this manuscript's audio summary by JACC Editor-in-Chief Dr. Valentin Fuster.



From the ^aDepartment of Medicine, Division of Cardiology, Department of Surgery, Division of Vascular Surgery, New York University School of Medicine, New York, New York; and the ^bDepartment of Medicine, Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine at UCLA, Los Angeles, California. Dr. Berger was supported, in part, by the National Heart, Lung, and Blood Institute of the National Institutes of Health (R01HL114978); receives research funding from AstraZeneca; has served on the executive committee for the EUCLID trial of antiplatelet therapy in patients with peripheral artery disease; and receives consulting fees from Janssen and Merck. Dr. Ladapo was supported by a K23 Career Development Award (K23 HL116787) from the National Heart, Lung, and Blood Institute (NHLBI) and the Robert Wood Johnson Foundation (74140).

Manuscript received February 14, 2017; accepted February 28, 2017.

ABBREVIATIONS AND ACRONYMS

ACEI = angiotensin-converting enzyme inhibitor

ARB = angiotensin receptor blocker

CAD = coronary artery disease

CI = confidence interval

NAMCS = National Ambulatory Medical Care Survey

NCHS = National Center for Health Statistics

NHAMCS = National Hospital Ambulatory Medical Care Survey

OR = odds ratio

PAD = peripheral artery disease

a coronary artery disease (CAD) risk equivalent (5). From a societal perspective, the consequences of PAD are significant (6).

SEE PAGE 2301

Given its high prevalence and poor prognosis, and shared risk factors with CAD, a major goal of PAD treatment includes risk factor modification and prevention of cardiovascular events (7). Guideline-directed therapy includes cardioprotective pharmacotherapies (i.e., antiplatelet therapy; statins; and blood pressure control, preferably with angiotensin-converting enzyme inhibitors [ACEIs] or angiotensin receptor blockers [ARBs]) and lifestyle counseling for healthy behaviors (i.e., diet, physical activity, and smoking cessation) (6,8). Adherence to pharmacological and lifestyle recommendations in PAD is uncertain. Several studies have shown that the use of proven cardioprotective medication for secondary prevention in patients with PAD significantly lags behind treatment for CAD (9–12). Previous analyses of PAD cohorts have been limited to populations enrolled in clinical trials, patients admitted to the hospital, patients undergoing lower extremity revascularization, or a single snapshot in time. Moreover, most studies of guideline adherence have focused on pharmacological therapies. Effective non-pharmacological therapies for PAD exist, including smoking cessation, regular physical activity, and diet counseling (6,8).

There is little known about national patterns of medication use and lifestyle counseling in patients with PAD in the United States; the extent to which trends may be attributable to changing population demographics, risk factors, and provider characteristics; or whether sex or racial/ethnic disparities exist in its use. Examining differences in this context is important because differences in the use (underuse) of medical therapies and lifestyle counseling could contribute to poorer cardiovascular health outcomes observed in certain groups. To answer these questions, we used nationally representative data to explore trends in medication use and lifestyle counseling in the United States among patients with PAD; investigate whether these trends may be attributable to shifts in population demographics, clinical risk factors, and provider characteristics; and evaluate whether sex or racial/ethnic disparities exist in such patients.

There is little known about national patterns of medication use and lifestyle counseling in patients with PAD in the United States; the extent to which trends may be attributable to changing population demographics, risk factors, and provider characteristics; or whether sex or racial/ethnic disparities exist in its use. Examining differences in this context is important because differences in the use (underuse) of medical therapies and lifestyle counseling could contribute to poorer cardiovascular health outcomes observed in certain groups. To answer these questions, we used nationally representative data to explore trends in medication use and lifestyle counseling in the United States among patients with PAD; investigate whether these trends may be attributable to shifts in population demographics, clinical risk factors, and provider characteristics; and evaluate whether sex or racial/ethnic disparities exist in such patients.

METHODS

DATA AND STUDY POPULATION. We analyzed data from the 2006 to 2013 National Ambulatory Medical

Care Survey (NAMCS) and National Hospital Ambulatory Medical Care Survey (NHAMCS), nationally representative surveys of adults seeking ambulatory care (13). We included all visits to office-based physicians and hospital-based outpatient clinics by adults (18 years of age or older; N = 418,889 adult visits). The National Center for Health Statistics (NCHS) and the Centers for Disease Control and Prevention conduct the NAMCS and NHAMCS annually on a nationally representative sample of visits to office-based physicians, hospital-based outpatient clinics, and emergency departments in the United States. For the NAMCS, each physician is randomly assigned to a 1-week reporting period during which a random sample of visits are surveyed systematically. Data are collected on patients' symptoms, comorbidity, and demographic characteristics; physicians' diagnoses; medications ordered or provided; and medical services provided. For the NHAMCS, a systematic random sample of patient visits in selected non-institutional general and short-stay hospitals are surveyed during a randomly assigned 4-week reporting period. The data collected on patient and provider characteristics are comparable to those collected in the NAMCS. Data on outpatient hospital departments and community health centers from the NHAMCS were unavailable in 2012 to 2013, but the majority of ambulatory care is performed in office-based visits and captured by the NAMCS (93% of visits during 2006 to 2011 occurred in the office rather than hospital outpatient departments, and 99% of office visits occurred outside of community health centers). However, we adjusted for the absence of these 2 care sites in our regression analyses and used the ratio of estimates derived from 2006 to 2011 with and without hospital outpatient/community health center visits to adjust 2012 to 2013 estimates of care provision and visit volume (14).

The NAMCS and NHAMCS intake materials allow physicians and staff to record up to 3 reasons for each visit and 3 diagnoses related to the visit, in addition to capturing several other major comorbid diagnoses (coded by NCHS staff using the International Classification of Diseases-Ninth Revision-Clinical Modification [ICD-9-CM]). From 2005 to 2013, the physician and hospital/outpatient clinic response rates in the NAMCS and NHAMCS ranged from 54% to 73% (except in 2012, when NAMCS response was 39%) and 80% to 95%, respectively, and item nonresponse rates were generally $\leq 5\%$ in both surveys.

STUDY POPULATION AND MEASURES. We identified patients with PAD on the basis of visit diagnoses using ICD-9 codes, including: 440.20-440.24, 440.29,

440.30-440.32, 440.4, 440.9, 443.81, 443.9, 445.02, and 785.4 (15). We used visit diagnoses and reasons for visit from our prior work to identify patients with other risk factors for adverse cardiovascular events, including hypertension, dyslipidemia, cigarette smoking, obesity, diabetes, chronic kidney disease, chronic obstructive pulmonary disease, CAD, and stroke (16,17). We were unable to validate our PAD diagnosis codes, but they were extracted from an American Heart Association Scientific Statement, and the NAMCS/NHAMCS uses rigorous methods to review and classify data reported on patient record forms.

We examined physician decision-making and treatment patterns for patients with PAD using: 1) Multum Lexicon drug codes and therapeutic drug categories and NCHS generic codes for antiplatelet agents (aspirin, clopidogrel, ticlopidine, and prasugrel), cilostazol, statins, ACEI/ARBs, and smoking cessation medications (nicotine replacement therapy, varenicline, or bupropion); and 2) physicians' reports of their provision of behavioral therapy (diet, weight loss, or exercise counseling) and smoking cessation therapy among smokers during the visit. A maximum of 8 medications could be recorded for visits between 2006 and 2011 and this increased to 10 medications in 2012 to 2013. We limited our accounting to the first 8 medications for each visit across all years for conformity but performed a sensitivity analysis in which up to 10 medications were assessed. This sensitivity analysis did not significantly change our results.

OTHER MEASURES. To examine factors associated with diagnostic and treatment patterns, we extracted information on demographic characteristics including age, sex, race/ethnicity, insurance (private, Medicare, Medicaid, self-pay/no-charge, and other/unknown), U.S. census region (Northeast, Midwest, South, and West), and urban or rural setting. We characterized patients as non-Hispanic white, non-Hispanic black, Hispanic, or other race. In subanalyses, physician specialty was divided into cardiologists and primary care physicians (general/family practice physicians and internists) because these physicians primarily manage PAD. Vascular surgeons were not included in these subanalyses because they were not identified in the dataset.

STATISTICAL ANALYSIS. We used summary statistics to estimate the prevalence of PAD and management patterns among patients with and without concurrent CAD. We estimated simple logistic regressions with year included as a continuous linear predictor to examine time trends. We also estimated


TABLE 1 U.S. Ambulatory Care Visits in Patients With PAD, by Demographic and Clinical Characteristics, 2006 to 2013

	All PAD			
	Unweighted Visits, n	Annual Weighted Visits, n	%	SE
All visits	1,982	3,781,624	100.0	0.0
Age, yrs				
<65	688	1,257,975	33.3	1.9
65-79	838	1,624,249	43.0	1.9
≥80	456	899,399	23.8	1.7
Sex				
Female	909	1,837,938	49.6	2.1
Male	1,073	1,907,686	50.4	2.1
Race/ethnicity				
Non-Hispanic white	1,142	2,124,148	56.2	2.6
Non-Hispanic black	241	383,805	10.1	1.8
Hispanic	152	290,825	7.7	1.1
Other/unknown	447	982,846	26.0	2.7
Insurance				
Private	422	953,056	25.2	2.0
Medicare	1,289	2,422,815	64.1	2.2
Medicaid	129	161,790	4.3	0.8
Other/unknown	81	171,348	4.5	1.0
Uninsured	61	72,614	1.9	0.6
U.S. region				
Northeast	373	698,106	18.5	2.4
Midwest	494	703,023	18.6	2.0
South	859	1,847,701	48.9	3.6
West	256	532,793	14.1	1.8
Setting				
Urban	1,791	3,349,979	88.6	2.3
Rural	191	431,644	11.4	2.3
Risk factor history				
Obesity	178	401,346	10.6	1.5
Smoker	416	771,949	20.4	1.7
COPD	215	452,067	12.0	1.3
Dyslipidemia	711	1,507,176	39.9	2.8
Diabetes	755	1,209,135	32.0	2.2
Hypertension	1,270	2,458,404	65.0	2.2
Chronic kidney disease	162	296,961	7.9	1.2
Comorbid cardiovascular diseases				
Coronary artery disease	425	912,113	24.1	1.9
Stroke	255	444,028	11.7	1.3

All analyses account for the complex sampling design of the NAMCS and NHAMCS.
CAD = coronary artery disease; COPD = chronic obstructive pulmonary disease; NAMCS = National Ambulatory Medical Care Survey; NHAMCS = National Hospital Ambulatory Medical Care Survey; PAD = peripheral artery disease; SE = standard error.

simple logistic regression models to examine differences in care among patients with PAD and CAD versus PAD alone, and among patients cared for by cardiologists versus primary care physicians. These models also adjusted for care site, based on whether visits occurred in a physician office, hospital outpatient department, or community health center. Multivariable logistic regression models adjusted for patients' clinical risk factors and demographic

FIGURE 1 Time Trends for Reported Medication Use and Lifestyle Counseling in Patients With PAD, 2006 to 2013



Data for medication use (A) include any antiplatelet therapy, statins, and ACEIs or ARBs. Data for lifestyle counseling (B) include physical activity or diet counseling and smoking cessation. ACEI = angiotensin-converting enzyme inhibitor; ARB = angiotensin receptor blocker; PAD = peripheral artery disease.


characteristics, insurance status, geographical region, setting (urban or rural), and care site. Analyses accounting for specialty were limited to the NAMCS because specialty information was unavailable in NHAMCS. To determine whether specific patient and provider characteristics accounted for any overall trends we observed, we constructed simple logistic regression models that assessed whether demographic factors or physician specialty changed in prevalence over the duration of our study period. All analyses accounted for the complex sampling design of the NAMCS and NHAMCS, and were performed using Stata version 14 (StataCorp, College Station, Texas) (18).

RESULTS

Over the 8-year period, the average annual number of ambulatory visits in the United States for PAD was 3,750,000. The estimated number of annual visits for PAD in the United States increased from 2.7 million (95% confidence interval [CI]: 1.9 to 3.5 million) in 2006 to 3.4 million (95% CI: 2.4 to 4.4 million) in 2013. The overall prevalence of a PAD diagnosis in adults in the dataset was 0.4%. Of all visits among patients with PAD, the prevalence of PAD with concomitant CAD was 24.1% and did not change significantly over time (Table 1). Patient age, sex, race/ethnicity distribution, geographic region distribution, and proportion with reported Medicare coverage also did not change significantly over time in patients with PAD, but the proportion of visits by women fell over time ($p < 0.01$). The prevalence of cardiologist visits did not change over time (22% in 2006 to 21% in 2013; $p = 0.88$ for trend).

PHARMACOLOGICAL THERAPY. The proportion of visits with reported use of antiplatelet therapy was 36.3% in 2006 to 2007 and 39.7% in 2012 to 2013, with no significant change over time ($p = 0.59$ for trend) (Figure 1A). Neither aspirin nor clopidogrel use changed over time. Concomitant use of dual antiplatelet therapy with aspirin plus clopidogrel was infrequent (7.3% in 2006 to 2007 to 7.1% in 2012 to 2013; $p = 0.38$ for trend). Visits by patients with PAD and CAD were more likely to report the patients being on antiplatelet therapy than patients with PAD alone (Figure 2). When stratified by coexistent CAD, visits for PAD without CAD reported a numerical increase in the use of any antiplatelet therapy (33.8% in 2006 to 2007 to 37.6% in 2012 to 2013) and aspirin (21% in 2006 to 2007 to 29.5% in 2012 to 2013) over time, though these trends were not statistically significant ($p = 0.72$ and $p = 0.38$, respectively). There was no change in antiplatelet therapy over time in patients with concomitant PAD and CAD (44.5% to 46.6%; $p = 0.43$ for trend).

The overall proportion of visits with reports of statin therapy was 35% (Table 2) and did not change significantly over time (30.5% in 2006 to 2007, to 38.8% in 2012 to 2013; $p = 0.18$ for trend) (Figure 1A). The most substantial increase in statin use was observed in patients with PAD alone, which trended toward significance (23.6% to 35.7%; $p = 0.057$ for trend). The proportion of visits with reports of ACEIs or ARBs was 31.1% and did not change significantly over time (32.9% in 2006 to 2007, to 29.7% in 2012 to 2013; $p = 0.51$ for trend). Use of cilostazol was noted in 5% of all visits and did not change over time ($p = 0.85$ for trend).



LIFESTYLE COUNSELING. The proportion of patient visits with reported exercise or diet counseling did not change over time (18% in 2006 to 2007, to 12.7% in 2012 to 2013; $p = 0.23$ for trend). In our visit sample, 20% of patients were current smokers, 51% of patients were past smokers, and smoking status was not reported in 29% of visits. Among smokers, smoking cessation counseling or pharmacotherapy was observed in 36.2% of visits, and this did not change over time (36.8% in 2006 to 2007, to 38.7% in 2012 to 2013; $p = 0.96$ for trend) (Table 2, Figure 1B).

PREDICTORS OF MEDICATION USE AND LIFESTYLE COUNSELING. We assessed predictors of medication use and lifestyle counseling by using data from the 1,982 surveys among patients with a PAD diagnosis in 2006 through 2013 (Online Table 1). Potential predictors included age, sex, race/ethnicity, insurance status, geographic region, setting, cardiovascular risk factors, comorbid cardiovascular disease, and time trend. The only significant variable associated with the use of antiplatelet therapy was concomitant CAD (odds ratio [OR]: 2.3; 95% CI: 1.6 to 3.4). Significant variables associated with statin use were dyslipidemia (OR: 3.3; 95% CI: 2.2 to 5.1) and concomitant CAD (OR: 1.6; 95% CI: 1.0 to 2.4). Use of ACEIs/ARBs was associated with dyslipidemia (OR: 1.6; 95% CI: 1.1 to 2.5), diabetes (OR: 1.6; 95% CI: 1.1 to 2.4), and hypertension (OR: 1.9; 95% CI: 1.2 to 3.0). We found no significant predictors for cilostazol use. Medication use did not differ over time in multivariate analyses.

Exercise or diet counseling was associated with obesity (OR: 3.5; 95% CI: 2.1 to 6.1) and hypertension


(OR: 1.6; 95% CI: 0.9 to 2.7) although the latter failed to reach statistical significance. Use of smoking cessation advice or medications was less likely in subjects ≥ 65 years of age (OR: 0.3; 95% CI: 0.1 to 0.7), non-whites (OR: 0.3; 95% CI: 0.1 to 0.9) and more likely in subjects with concomitant CAD (OR: 3.5; 95% CI: 1.6 to 7.4). Lifestyle counseling did not differ over time in multivariate analyses (Online Table 1).

PHYSICIAN TYPE. Because quality of care can be dependent by physician type (19), we sought to investigate medication use and lifestyle counseling by practicing physician. Specialty data are not

TABLE 2 Prevalence of Medication Use or Lifestyle Counseling in Patients With PAD Seeing Physicians in U.S. Ambulatory Care Visits, 2006 to 2013

	All PAD, Years 2006-2013			
	Unweighted Visits, n	Annual Weighted Visits, n	%	SE
Medical therapy				
Any antiplatelet therapy	633	1,400,619	37.8	2.6
Aspirin	497	1,059,044	28.6	2.2
Clopidogrel	287	699,383	18.1	1.9
Statin	557	1,297,320	35.0	2.4
ACEI/ARB	482	1,152,083	31.1	2.0
Cilostazol	71	186,216	5.0	1.0
Lifestyle counseling				
Exercise or diet counseling	299	745,081	20.1	2.2
Smoking cessation	146	274,752	36.3	3.9

All analyses account for the complex sampling design of the NAMCS and NHAMCS.
ACEI = angiotensin converting enzyme inhibitor; ARB = angiotensin receptor blocker; other abbreviations as in Table 1.



available in NHAMCS (hospital outpatient), and thus a separate sensitivity analysis was performed with NAMCS (physician office) data. Cardiologists and primary care physicians were the most likely groups taking care of these patients. As noted in **Figure 3**, PAD visits with a cardiologist were more likely to be on antiplatelet therapy, statins, ACEIs or ARBs, and more frequently counselled for smoking cessation ($p < 0.05$ for all). After multivariate analysis, antiplatelet therapy was significantly more likely to be used in PAD when seen by a cardiologist (OR: 1.8; 95% CI: 1.1 to 2.9). Use of statins, ACEIs or ARBs, and lifestyle counseling were not significantly different between groups.


DISCUSSION

PAD is an increasingly common medical problem, and numerous clinical studies and guidelines have focused on its optimal treatment. To investigate the use of clinical guidelines, it is important to monitor physician practice patterns. The NAMCS and the NHAMCS provide unique insights into the clinical management of PAD in the United States over time. In the present study, we examined a representative sample of visits to physicians in the United States between 2006 and 2013 among patients with PAD. The proportion of visits reporting use of medical therapy and lifestyle counseling was low and appears suboptimal in consideration of the substantial benefit of secondary cardiovascular prevention and lifestyle counseling (**Central Illustration**). These

findings suggest that a considerable proportion of patients with PAD disease remain at increased risk for adverse outcomes.

Our findings clearly show the underuse of cardiovascular prevention medication in patients with PAD. It has been well documented that antiplatelet therapy and statins significantly reduce incident cardiovascular events in PAD (6,7). Although use of antiplatelet therapy and statins increased from 2006 to 2013, the magnitude of this increase was less than expected, and this increase was not significant after adjusting for demographics and clinical characteristics of the patients. In multivariate analysis, sex and race/ethnicity of the patient did not play a role in the use of secondary prevention and lifestyle counselling. Although we are unable to determine the reason for the underuse of secondary prevention and lifestyle counselling, interventions to optimize secondary prevention, such as physician note “checklists” or other systematic prescription programs, would likely improve medication use and lifestyle counselling (20,21).

Although some, but not all, prior studies have reported comparatively higher rates of secondary prevention in patients with PAD, many of these studies focused on selected patients whose use of secondary prevention might be higher than that in the physician office setting assessed here (9,11,22). Studies of hospitalized patients have shown higher rates, likely due to the focused clinical attention that these patients receive. Hospitalized patients, however, constitute a minority of all patients with PAD. Practices in community settings may be more likely to reflect the



public health impact of lifestyle recommendations and secondary prevention efforts.

Similar to prior studies, we demonstrate that secondary prevention use in PAD was not uniform across patient subpopulations. In particular, secondary prevention was less likely to be reported in patients with PAD alone (without concomitant CAD). Although patients with PAD and CAD represent a particularly high-risk group, patients with PAD alone are still at heightened risk for cardiovascular and limb events, and impairment in quality of life. The greater use of secondary prevention medication in patients with PAD and CAD may indicate a higher use of medication in patients with polyvascular disease or the importance of secondary prevention in CAD irrespective of PAD. These data suggest important barriers to the uniform and widespread adoption of secondary prevention in PAD. The underuse of lifestyle recommendations and cardiovascular prevention in this population is a major concern and supports the need for better awareness and education for the vascular community.

STUDY LIMITATIONS. The NAMCS and NHAMCS provide a limited amount of clinical information on each patient visit; there are no data on the severity of PAD nor could we characterize the symptom status of the patient. Our estimates of medication use could also be lower than expected since underreporting of medication use and lifestyle counseling may have occurred, in particular, aspirin and diet counseling. The NAMCS and NHAMCS do not capture the longitudinal experience of the individual patients, and because the unit of measurement is by patient visit, frequent users of care may be overrepresented. Finally, we are unable to

determine specific reasons that patients are not taking medications, such as adverse drug reactions or patient preference. Thus, determining appropriateness of care is difficult. In addition, it may have been more appropriate to compare our patients with PAD and concurrent CAD to patients with CAD alone, but our analysis did not incorporate the latter cohort.

CONCLUSIONS

This analysis suggests that use of secondary prevention and lifestyle counseling in patients with PAD has not become a widely disseminated practice in the United States, a finding consistent with a global pattern (9,11,12). Our study identifies important targets for immediate improvement of health care outcomes in patients with PAD. New health care system strategies are required to ensure adequate resource utilization in patients with PAD. Although much attention is focused on novel therapies in PAD, a refocus on established therapies and healthy behaviors is clearly needed. Attempts to increase patient and physician awareness of the benefits of lifestyle recommendations and secondary prevention may be necessary. In addition, systems of chronic disease management in which the use of nurses, other health care providers, or information systems complements the role of physicians also may be helpful. Finally, efforts to monitor the prevention practices taking care of PAD patients may provide new incentives for quality care. The personal, societal, and financial burdens of preventable deterioration of patients with PAD suggest that a substantial investment in such

strategies may be warranted. Future research aimed at improving secondary prevention and lifestyle counseling in patients with PAD is certainly needed.

ADDRESS FOR CORRESPONDENCE: Dr. Jeffrey S. Berger, Department of Medicine, Division of Cardiology, Department of Surgery, Division of Vascular Surgery, New York University School of Medicine, 530 First Avenue, SKR-9R, New York, New York 10016. E-mail: Jeffrey.berger@nyumc.org. OR Dr. Joseph A. Ladapo, Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine at UCLA, 10833 LeConte Avenue, #12138, Los Angeles, California 90095. E-mail: JLadapo@mednet.ucla.edu.

PERSPECTIVES

COMPETENCY IN SYSTEMS-BASED PRACTICE: Guideline-recommended measures for lifestyle interventions to reduce the risk of cardiovascular events are utilized in fewer than one-half of the patients with PAD managed in ambulatory care practice.

TRANSLATIONAL OUTLOOK: More work is needed to develop systems of care that promulgate evidence-based interventions to improve clinical outcomes in this high-risk patient population.

REFERENCES

1. Fowkes FG, Rudan D, Rudan I, et al. Comparison of global estimates of prevalence and risk factors for peripheral artery disease in 2000 and 2010: a systematic review and analysis. *Lancet* 2013;382:1329-40.
2. Savji N, Rockman CB, Skolnick AH, et al. Association between advanced age and vascular disease in different arterial territories: a population database of over 3.6 million subjects. *J Am Coll Cardiol* 2013;61:1736-43.
3. Mozaffarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statistics-2016 update: a report from the American Heart Association. *Circulation* 2016;133:e38-360.
4. Criqui MH, Aboyans V. Epidemiology of peripheral artery disease. *Circ Res* 2015;116:1509-26.
5. Subherwal S, Patel MR, Kober L, et al. Peripheral artery disease is a coronary heart disease risk equivalent among both men and women: results from a nationwide study. *Eur J Prev Cardiol* 2015;22:317-25.
6. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2017;69:e71-126.
7. Berger JS, Hiatt WR. Medical therapy in peripheral artery disease. *Circulation* 2012;126:491-500.
8. Hivert MF, Arena R, Forman DE, et al. Medical training to achieve competency in lifestyle counseling: an essential foundation for prevention and treatment of cardiovascular diseases and other chronic medical conditions: a scientific statement from the American Heart Association. *Circulation* 2016;134:e308-27.
9. Cacoub PP, Abola MT, Baumgartner I, et al. Cardiovascular risk factor control and outcomes in peripheral artery disease patients in the Reduction of Atherothrombosis for Continued Health (REACH) registry. *Atherosclerosis* 2009;204:e86-92.
10. Pande RL, Perlstein TS, Beckman JA, Creager MA. Secondary prevention and mortality in peripheral artery disease: National Health and Nutrition Examination Study, 1999 to 2004. *Circulation* 2011;124:17-23.
11. Subherwal S, Patel MR, Kober L, et al. Missed opportunities despite improvement in use of cardioprotective medications among patients with lower-extremity peripheral artery disease, underuse remains. *Circulation* 2012;126:1345-54.
12. Cacoub PP, Zeymer U, Limbourg T, et al. Effects of adherence to guidelines for the control of major cardiovascular risk factors on outcomes in the Reduction of Atherothrombosis for Continued Health (REACH) Registry Europe. *Heart* 2011;97:660-7.
13. National Center for Health Statistics (U.S.). Ambulatory Health Care Data: NAMCS and NHAMCS Description. 2013. Available at: http://www.cdc.gov/nchs/ahcd/ahcd_questionnaires.htm. Accessed June 4, 2013.
14. National Center for Health Statistics (U.S.). 2009 NAMCS Public-Use Data File Documentation. Available at: http://www.cdc.gov/nchs/ahcd/ahcd_questionnaires.htm. Accessed June 4, 2013.
15. Hirsch AT, Allison MA, Gomes AS, et al. A call to action: women and peripheral artery disease: a scientific statement from the American Heart Association. *Circulation* 2012;125:1449-72.
16. Ladapo JA, Blecker S, Douglas PS. Physician decision making and trends in the use of cardiac stress testing in the United States: an analysis of repeated cross-sectional data. *Ann Intern Med* 2014;161:482-90.
17. Sigmund AE, Stevens ER, Blitz JD, Ladapo JA. Use of preoperative testing and physicians' response to professional society guidance. *JAMA Intern Med* 2015;175:1352-9.
18. National Center for Health Statistics (U.S.). Ambulatory Health Care Data: NAMCS and NHAMCS, Reliability of Estimates. 2013. Available at: http://www.cdc.gov/nchs/ahcd/ahcd_estimation_reliability.htm. Accessed June 4, 2013.
19. Donohoe MT. Comparing generalist and specialty care: discrepancies, deficiencies, and excesses. *Arch Intern Med* 1998;158:1596-608.
20. Cebul RD, Love TE, Jain AK, Hebert CJ. Electronic health records and quality of diabetes care. *N Engl J Med* 2011;365:825-33.
21. Jepson RG, Harris FM, Platt S, Tannahill C. The effectiveness of interventions to change six health behaviours: a review of reviews. *BMC Public Health* 2010;10:538.
22. Armstrong EJ, Chen DC, Westin GG, et al. Adherence to guideline-recommended therapy is associated with decreased major adverse cardiovascular events and major adverse limb events among patients with peripheral arterial disease. *J Am Heart Assoc* 2014;3:e000697.

KEY WORDS cardiovascular prevention, epidemiology, lifestyle, PAD

APPENDIX For a supplemental table, please see the online version of this article.